

<https://helda.helsinki.fi>

---

## Evaluation of Tail Lesions of Finishing Pigs at the Slaughterhouse : Associations With Herd-Level Observations

Heinonen, Mari

2021-07-15

---

Heinonen , M , Välimäki , E , Laakkonen , A-M , Toppari , I , Vugts , J , Fabrega , E & Valros , A 2021 , ' Evaluation of Tail Lesions of Finishing Pigs at the Slaughterhouse : Associations With Herd-Level Observations ' , Frontiers in Veterinary Science , vol. 8 , 650590 . <https://doi.org/10.3389/fvets.2021.650590>

---

<http://hdl.handle.net/10138/333101>

<https://doi.org/10.3389/fvets.2021.650590>

---

cc\_by

publishedVersion

---

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.



# Evaluation of Tail Lesions of Finishing Pigs at the Slaughterhouse: Associations With Herd-Level Observations

Mari Heinonen <sup>1\*</sup>, Elina Välimäki <sup>2</sup>, Anne-Maija Laakkonen <sup>1</sup>, Ina Toppari <sup>3</sup>, Johannes Vugts <sup>2</sup>, Emma Fàbrega <sup>4</sup> and Anna Valros <sup>1</sup>

<sup>1</sup> Department of Production Animal Medicine and Research Centre for Animal Welfare, Faculty of Veterinary Medicine, University of Helsinki, Helsinki, Finland, <sup>2</sup> Animal Sourcing, HKScan Finland Oy, Forssa, Finland, <sup>3</sup> Animal Health ETT, Seinäjoki, Finland, <sup>4</sup> Animal Welfare Program, Institut de Recerca i Tecnologia Alimentària, Monells, Spain

## OPEN ACCESS

Edited by:  
Stephanie Torrey,  
Independent Researcher,  
Guelph, Canada

Reviewed by:  
Nienke Van Staaveren,  
University of Guelph, Canada  
Jen-Yun Chou,  
University of Pennsylvania,  
United States

\*Correspondence:  
Mari Heinonen  
mari.heinonen@helsinki.fi

Specialty section:  
This article was submitted to  
Animal Behavior and Welfare,  
a section of the journal  
Frontiers in Veterinary Science

Received: 07 January 2021

Accepted: 13 April 2021

Published: 15 July 2021

Citation:  
Heinonen M, Välimäki E,  
Laakkonen A-M, Toppari I, Vugts J,  
Fàbrega E and Valros A (2021)  
Evaluation of Tail Lesions of Finishing  
Pigs at the Slaughterhouse:  
Associations With Herd-Level  
Observations.  
Front. Vet. Sci. 8:650590.  
doi: 10.3389/fvets.2021.650590

The prevalence of tail lesions evaluated at the slaughterhouse varies considerably between herds. These lesions result mainly from tail biting, a harmful behavior with multifactorial origin. This study sought to investigate if a batchwise inspection of tails at slaughterhouse could be a useful method to estimate the animal welfare situation in finishing pig herds, and if so, what type and detail of tail scoring such an inspection should utilize. We investigated the distribution of different types of tail lesions and how well their scoring at slaughterhouse was associated with the situation recorded on-farm by a veterinarian as part of routine herd health visits. We also wanted to determine if animal welfare-related herd-level parameters, recorded by herd veterinarians during herd health visits, are associated with tail scoring at the slaughterhouse. A total of 10,517 pig tails from 84 herds were scored for this study. Herd data were collected from the national health classification register for pig farms in Finland and also included annual herd production quality data collected by the slaughterhouse. The scores of the tails varied considerably between the herds. On average, 48.1% (sd D 19.3) of the tails with an average length of 30.4 cm (sd D 2.7) were fully intact, 37.3% (13.9) had healed (length D 26.4, sd D 5.1 cm), 12.4% (9.0) (length D 28.9, sd D 4.3 cm) had minor acute wounds, and 2.3% (2.1) (length D 24.2, sd D 6.0 cm) had major acute wounds. Proportions of different tail lesions at slaughterhouse were associated with or tended to be associated with the following herd-level parameters in regression models: use of wood as enrichment ( $p < 0.1$ ), one health parameter (leg problems other than arthritis,  $p < 0.05$ ), and long-term animal welfare estimate (annual mortality,  $p < 0.05$ ). Detailed tail evaluation at the slaughterhouse shows potential in estimating the tail lesions and long-term welfare level on the farm. By recording only one type of tail condition (such as tails with major acute lesions) at the slaughterhouse, it is not possible to estimate the total tail lesion situation in the herds before slaughter. A more detailed scoring similar to the one used in this trial is recommended.

**Keywords:** tail biting, pig welfare, herd health, tail scoring, abattoir, harmful behavior

## INTRODUCTION

Tail biting (TB) is a major welfare concern in pig production (1) and also leads to large economic losses (2). Tail biting results in tail lesions (TL) and is related to several other problems, such as infection in the bitten pig (3), stress and pain in pigs (4, 5), carcass condemnations (6–9), and reduced daily gain (10). TB is a harmful behavior and has a multifactorial origin. Both suboptimal environmental, health and management-related factors, and certain characteristics of individual pigs are risk factors for TB (11). The magnitude of the number of possible risk factors is illustrated in the study of Taylor et al. (12), who collected altogether 83 risk factors for their husbandry advisory. It is important to be aware of risk factors for TB, these parameters have an important effect on the welfare of pigs. Thus, tail lesions in the herds could be used as a simple measure of animal welfare, either at herd level before slaughter or at slaughterhouse during meat inspection. Previous studies have combined the estimation of tail lesions at the slaughterhouse and herd-level data (4). Meat inspection of pigs has been suggested as a welfare surveillance tool (8), and other research has shown potential to use TL scoring as indicators of pig welfare on farms (5–13).

The prevalence of TL varies considerably in scientific studies and has been estimated to be 2–4 fold higher in pigs with undocked tails than those with docked tails (9). Sampling methods and definitions for TL differ and therefore studies often cannot be compared. The results differ especially when the tails were evaluated in the herd before slaughter or the data were collected from routine meat inspection data. In addition, it has been common in slaughterhouse studies to include mild lesions in the same score with healed lesions (18, 20–24) or healed lesions have not been mentioned separately in the scoring definitions (9, 25). Only a few studies report healed tails separately from other tail lesions (26). It is also not common to report the variability of different lesions in the slaughterhouses. For example, the results from some studies show that severe tail lesions can be found in an average of 1.2–1.9% of the tails scored at the slaughterhouse; the minimum in all studies was 0% and the maximum varied between 7 and 22.9% (8, 23). We consider both of these parameters very important. Healed tails can give an indication of when the tail damage occurred and knowledge on the variability of different TLs in the batches may help in preventing different types of TB in the herds (7).

Our aim was to investigate if a batchwise, thorough inspection of tails at the slaughterhouse could be a useful method to assess the animal welfare situation in finishing pig herds, and if so, what type and detail of tail scoring such an inspection should utilize. To achieve this, we investigated (a) how the distribution of different types of intact and damaged tails vary between farms, (b) how well TL scoring at the slaughterhouse is associated with the herd-level situation of tails scored during a single health visit, and (c) how long term production parameters (annual mortality, annual total, and partial carcass condemnations) are associated with tail scores in the slaughterhouse. In addition, to further evaluate the value of TL scoring at the slaughterhouse as a proxy measure of on-farm conditions: (d) we tested if certain relevant, and available, risk factors for tail biting (mainly related

to health and use of enrichment), recorded by herd veterinarians during herd health visits, are associated with tail scores in the slaughterhouse. We hypothesized that by recording only the severely bitten pigs at the slaughterhouse we could estimate the total TL level in the herd, both in the long and short term. We also hypothesized that certain herd-level parameters, such as animal health and management features, are associated with the level of tail damage scored at the slaughterhouse.

## MATERIALS AND METHODS

### Tail Scoring at the Slaughterhouse

The data collection from undocked finishing pigs has been described in detail by Valros et al. (27). Here we describe only the main points of scoring. The data were collected at a large Finnish slaughterhouse over 5 consecutive days in June 2019. Most of the carcass tails handled at the slaughterhouse were scored. In this study, we included only the tails from the herds that had sent at least 50 pigs to slaughter during the data collection period of 5 days, leaving a sample of 10,517 tail scores from 84 herds (see more in data handling and statistical analyses). We wanted to ensure better data quality by including only herds large enough to yield data not only from a few occasional animals but also from a large number of pigs. The carcasses were hung on gambrels ensuring the identification of the farm or origin. Carcasses were identified by the herd tattoo and individual slaughter data was recorded per pig via the gambrel identification system. Pork producers selling their finishers to slaughter during that week did not know about the study. This dataset includes a subsample of the carcasses reported in another article (26) (where the average carcass weight of the pigs was 89.7 kg and 138 kg).

Pictures and live observations were used to train the researchers to harmonize the scorings before the actual data collection at the slaughterhouse line. The scoring point was situated after singeing, whipping, bung drilling, and chest opening. Altogether six researchers, two at a time, were responsible for the scorings, which took about 7–8 s per tail due to slaughterhouse line speed. One of the two main scorers was present during all scoring sessions. The scorers were standing on a platform high enough for them to palpate and measure each tail easily. One researcher measured the tail of the carcass with a 50 cm long ruler and assessed the possible tail lesion while the other recorded the observations. With the carcasses hanging upside down, the end of the ruler was placed on the dorsal side of the tail and firmly pushed toward the base of the tail. The tail was then manually extended against the ruler. During scoring, the observers consulted each other actively in case of questionable scorings.

The scoring system used included elements from the systems developed by The FareWellDock-consortium (17) and those suggested by the pig welfare subgroup of the EU animal welfare platform (28). The preliminary system was amended to be usable at the slaughterhouse line after the pilot visit testing. A tail with multiple lesions was scored according to the most severe lesion. Tails scored as healed were also checked for acute damage. Finally, the tails were classified as intact, healed, with an acute

**TABLE 1 |** Tail scoring system used at the slaughter line after scalding of the carcass and the mean percentage and standard deviation of one delivery group of finishing pigs from 84 herds (10,517 carcasses) according to their tail lesion scores and tail length in centimeters.

Tail score	Definition	Mean (sd) percentage of tails	Tail length, mean, cm (sd)
Intact	The tail is fully intact, the end is rounded, and slightly attenuated	48.1% (19.3)	30.4 (2.7)
Healed	The tail is clearly shortened; the tail end is scarred, of abnormal shape or too thick to be intact. The skin is totally healed (no scab, wound, or missing tissue)	37.3% (13.9)	26.4 (5.1)
Minor acute wound	The tail has missing tissue, which has not yet fully healed; uneven dents in the skin; or a part of the tail is missing. Wound is > 0 cm but < 2 cm in diameter or length	12.4% (9.0)	28.9 (4.3)
Major acute wound	The tail has missing tissue, which has not yet fully healed; uneven dents in the skin; or a part of the tail is missing. Wound is ≥ 2 cm in diameter or length	2.3% (2.1)	24.2 (6.0)

wounds, and with major acute wounds (Table 1). Additional details are available in Valros et al. (6).

## Data Collection From the Herds Through the National Herd Health Register, Sikava

Data concerning the finishing pigs in 84 herds were collected from the national health classification register for pig farms in Finland, Sikava ([www.sikava.fi](http://www.sikava.fi)). The Sikava system has been ISO9001:2015 certified since 2014. The national level of Sikava fulfills the national food quality schemes criteria as described in Article 16 of Regulation (EC) No 1305/2013 and was recognized by Finnish Food Safety Authority EVIRA in 2013. At first, the data were checked and yielded a data set of 14,382 tails scored out of a total of 14,433 original scores. This classification system covers 90% of Finnish swine farms and over 95% of the pork production in the country. All herds of this study fulfilled the national level requirements of Sikava.

Detailed information of the health classification system can be found at [www.sikava.fi](http://www.sikava.fi). Here, we present only the most important features of the system relevant for this study. The herds that have not this study. national level of Sikava make a healthcare agreement with a veterinarian who has participated in a course organized by Sikava. The farm must invite the herd health veterinarian to visit the herd four to six times a year or at least once in every batch in all in-house all-out finishing units and the veterinarians need to document their observations in an internet-based database with periodic evaluations. The visit interval is controlled by Sikava and the delayed visits affect animal trade. The farm must be at a national level of Sikava before slaughter of the pigs.

In this study, we included observations about pig health, behavior and use of enrichment for the pigs recorded by the veterinarian during the herd health visit according to the scoring system acquired by Sikava (Table 2). The results of a single herd health visit record from each herd were extracted from the Sikava database. We used the latest visit documented by the herd health veterinarian before the data collection at the slaughterhouse. To ensure the visit included observations on the pigs assessed for damage at the slaughterhouse, only herds that had been visited between > 1 week and < 3 months before the first sampling day were included. This time period was an average of 36.8 (sd 21.8) days in our 84 study herds.

The slaughterhouse continuously collects the following information on parameters for follow-up of production quality: injuries/infections and “abscesses.” Secondly, a total score and saves this data in Sikava database: annual mortality and annual percentage of carcasses with total carcass condemnation.

and annual percentage of carcasses with partial carcass condemnation. The herd owners and health care veterinarian can see these parameters and use them as long-term animal health and welfare estimates for the farms and follow-up of production quality. In this project, we refer them as production parameters. They were available for 79 herds of this study for the period from July 2018 to June 2019.

## Data Handling and Statistical Analyses

All statistical analyses were performed using IBM SPSS statistics version 25. The unit of interest was the herd.

First, the data were checked and yielded a data set of 14,382 tails scored out of a total of 14,433 original scores. After including only the herds that had sent at least 50 pigs to slaughter during the study week and from which the herd health visit data were available (84 herds), altogether 10,517 (mean of 215.2 [sd 83.0] tails per herd) were included in the dataset of this study.

Continuous variables were assessed for normality visually and using the Shapiro–Wilk test. The correlation between the percentages of different tail scorings at the slaughterhouse was calculated with Pearson correlation. Differences between farms within different tail lesion percentage class, based on data collected by the veterinarian during the herd health visit (tail lesions below 1 vs. 1–10%) in the tail scoring at the slaughterhouse was evaluated using separate tests for each tail lesion outcome. Similarly, the difference between farms within the different percentage of intact tails-classes based on the herd health visit (80%, > 80%, and > 95% intact tails) in their percentage of tail outcomes at the slaughterhouse was tested using one-way ANOVA tests, followed by Bonferroni-correction for pairwise tests. After calculation of the descriptive preliminary univariable statistical analyses were performed to find the associations between the predictor variables (code from Sikava database, see Table 2) and the four different tail lesion outcomes one at a time (percentage of tail scores: intact, healed, minor acute wounds, and major acute wounds). Two new variables were formed by combining the evaluations performed by the veterinarians during the herd health visits. First, the “skin problems” represented the sum of the scores of the variables “skin lesions,” “injuries/infections,” and “abscesses.” Secondly, a total score was calculated as the sum of the scores given to all 11 symptoms (scored each as 0, 1, or 2 and thus yielding a



different tail outcomes in the slaughterhouse (intact tails, healed tails, tails with minor wounds, and tails with major wounds) were tested separately with Pearson correlations (total and partial carcass condemnations) and Spearman rank correlation (annual mortality, which was non-normally distributed).

RESULTS

Descriptive Statistics of Tail Scoring at the Slaughterhouse

Out of the 84 herds in this study, 69 (82.1%) were finishing units growing feeder pigs from a weight of 80 kg until slaughter

and 15 (17.9%) were farrow-to-finish herds. The production type did not affect the prevalence of the different tail scorings at the slaughterhouse ( $p > 0.05$  for all). The study herds had room for an average of 1,407.0 (SD 1,460.8) finishing pigs, and the average inter-visit mortality was 1.3% (SD 0.9).

On average, the mean percentage of fully intact tails at the slaughterhouse was 48.1% (Table 1 for the tail classification results). The percentage of intact tails correlated negatively with the percentage of healed tails and tails with minor and major wounds (see Table 3 for all correlations between different tail classifications). Figure 1 shows the frequency distributions of tail classifications in the herds. The study herds had varying combinations of the percentages of tail findings (Figure 2).

TABLE 3 | Correlation of percentages of herd-level tail evaluations from 84 Finnish herds.

Percentage of tails	Healed	With minor wounds	With major wounds
Intact	0.858**	0.706**	0.539**
Healed		0.258*	0.270*
With minor wounds			0.522**

Pearson correlation, \*\* $p < 0.01$  and \* $p < 0.05$ . Undocked tails of finishing pigs (n = 10,517) were evaluated at the slaughterhouse after scalding.

Descriptive Statistics of Sikava Herd Health Visit Recordings and Their Association With Slaughterhouse Tail Scoring

During the one herd health visit, the herd veterinarians scored TLs in 21 (25%) herds in none or single animals. In addition, 58 (69%), 5 (6.0%), and 0 herds were given a score of 1 (1–5% of the pigs), 2 (6–10% of the pigs), or 3 (10% of the pigs), respectively. Later in the analyses, the herds were divided in the following two groups: 21 herds with no or single animals with bitten tails







and 63 herds with TL in 1–10% of the pigs. Figure 3 shows the percentage of pigs with bitten tails to be 1% (score 0) compared to the different tail scores at slaughterhouse divided according to the evaluation recorded in the same herds during the herd health visit. In the herds where the veterinarian had estimated the percentage of the tails were evaluated at the slaughterhouse

**FIGURE 4 |** The average percentage of undocked tails of finishing pigs per herd (n = 84) scored at the slaughterhouse in four categories (intact, healed, with minor acute wounds, and with major acute wounds) and divided according to the evaluation of the percentage of intact tails recorded during the herd health visits (> 95%, > 80%, or < 80%). Within tail score comparisons, different letters represent statistical significance: ab  $p < 0.01$ , cd  $p < 0.05$ .

as intact and a lower percentage as tails with major acute wounds.

The veterinarians scored the percentage of intact tails in finishers during the herd health visit as follows: in 43 herds (51.2%) > 95% of the tails were intact, in 23 herds (27.4%) 80% of the tails were intact, in 16 herds (19.0%) 70% of the tails were intact, and one herd (1.2%) had 70% intact tails. Figure 4 shows the different tail scores at slaughterhouse divided according to the evaluation of the percentage of intact tails by the herd health veterinarian during the herd health visit: when the herd veterinarian had evaluated the percentage of intact tails to be > 95% in the herd compared to the figure of 80%, a higher percentage of the tails were scored at the slaughterhouse as intact and a lower percentage as tails with healed, minor acute or major acute wounds.

Descriptive herd data (84 herds) that were used in statistical analyses are shown in Table 4 (enrichment materials used) and Table 5 (disease symptoms evaluated by the veterinarian). The herd veterinarian had scored the explorative behavior to give material to be Good in 46 (54.8%), Satisfactory in 26 (30.9%), and Poor in 11 herds (13.1%). Similarly, the reaction to the examination was scored to be Good, Satisfactory, and Poor in 79 (94.9%), 4 (4.8%), and 0 herds, respectively.

### Regression Models for Outcomes Intact Tails, Healed Tails, Tails With Minor Wounds, and Tails With Major Wounds

The binomial regression model for percentage of intact tails ( $R^2 = 0.175$ ) included the following parameters: wood as enrichment material (b = 2.0, df = 1, F = 3.850, and  $p = 0.05$ ), with the association being positive. The binomial regression model for percentage of tails with major acute wounds ( $R^2 = 0.238$ ) included the following parameters: wood as enrichment material (b = 8.939, and  $p = 0.04$ ), explorative behavior (b = 2.845, and  $p = 0.1$ ), and the

**TABLE 4 |** Information recorded by veterinarian during one herd health visit on the use of different materials used as environmental enrichment in 84 study herds growing finishing pigs.

Material*	0 D No	1 D Yes
Toy	16 (19.0%)	68 (81.0%)
Straw	21 (25.0%)	63 (75.0%)
Sawdust	27 (32.1%)	57 (67.8%)
Wood	56 (66.7%)	28 (33.3%)
Paper	58 (69.0%)	26 (31.0%)
Peat	64 (76.1%)	20 (23.8%)
Hay	65 (77.4%)	19 (22.6%)
Wood chips	83 (98.8%)	1 (1.2%)
Other	82 (97.6%)	2 (2.4%)

\*The farms may have used more than one material.

where the veterinarian had recorded leg problems other than arthritis, the percentage of intact tails was higher if wood was used as enrichment compared with the situation when wood was not used (Figure 5).


The binomial regression model for healed tails ( $R^2 = 0.061$ ) included one parameter "leg problems other than arthritis" (b = 1, F = 5.293, and  $p = 0.02$ ). The herds with leg problems other than arthritis had an average of 40.5% (s.d. = 2.0) healed tails compared with no such symptom recorded (33.7% s.d. = 2.1). Similarly, the regression model for percentage of tails with minor wounds ( $R^2 = 0.045$ ) contained one parameter, inter-visit mortality (b = 2.0, df = 1, F = 3.850, and  $p = 0.05$ ), with the association being positive.

The binomial regression model for percentage of tails with major acute wounds ( $R^2 = 0.238$ ) included the following parameters: wood as enrichment material (b = 8.939, and  $p = 0.04$ ), explorative behavior (b = 2.845, and  $p = 0.1$ ), and the



TABLE 5 | The presence of different symptoms in the finishing pigs (number and percentage of herds out of 84) recorded by herd veterinarians during one herd health visit.

Symptom	0 D None or single animals	1 D Some, 1–5% of the pigs	2 D Several, 6–10% of the pigs	3 D Plenty, > 10% of the pigs
Arthritis	10 (11.9%)	68 (81.0%)	6 (7.1%)	0
Claw injury	68 (81.0%)	16 (19.0%)	0	0
Leg problem other than arthritis	40 (47.6%)	44 (52.4%)	0	0
Coughing	73 (86.9%)	8 (9.5%)	3 (3.6%)	0
Sneezing	77 (91.7%)	7 (8.3%)	0	0
Diarrhea	66 (78.6%)	18 (21.4%)	0	0
Skin injury or infection	44 (52.4%)	40 (47.6%)	0	0
Abscesses	50 (59.5%)	34 (40.5%)	0	0
Central nervous system symptoms	79 (94.0%)	5 (6.0%)	0	0
Runts	36 (42.8%)	48 (57.1%)	0	0
Hernia	10 (11.9%)	71 (84.5%)	3 (3.6%)	0



interaction “Wood as enrichment Explorative behavior” ( $r_s = 0.21, p = 0.07$ ), and annual mortality ( $r_s = 0.19, p = 0.09$ ) of the herd. The percentage of healed tails in the slaughterhouse data correlated positively with annual mortality ( $r_s = 0.26, p = 0.02$ ) but not with the measures of condemnations. The percentage of tails with minor wounds in slaughterhouse correlated positively both with annual total carcass condemnations ( $r_s = 0.30, p = 0.007$ ) and with annual partial carcass condemnations ( $r_s = 0.26, p = 0.02$ ) but not with annual mortality. The percentage of tails with major wounds scored in the slaughterhouse did not correlate with any of the parameters of production parameters ( $r_s < 0.1$  for all).

### Production Parameters and Their Association With Slaughterhouse Tail Scoring

The production parameters (long-term animal health and welfare estimates of the herds) revealed an average annual mortality of 1.9% (sd 1.88), an annual total carcass condemnation of 0.27% (0.17), and an annual partial carcass condemnation of 6.0%.

The percentage of intact tails in the slaughterhouse data tended to correlate negatively with annual total carcass condemnations ( $r_s = 0.21, p = 0.06$ ), annual partial carcass scores (intact, healed, minor wounds, and major wounds) ( $r_s = 0.19, p = 0.09$ ), and annual mortality ( $r_s = 0.26, p = 0.02$ ) but not with the measures of condemnations. The percentage of tails with minor wounds in slaughterhouse correlated positively both with annual total carcass condemnations ( $r_s = 0.30, p = 0.007$ ) and with annual partial carcass condemnations ( $r_s = 0.26, p = 0.02$ ) but not with annual mortality. The percentage of tails with major wounds scored in the slaughterhouse did not correlate with any of the parameters of production parameters ( $r_s < 0.1$  for all).

### DISCUSSION

Our study revealed that the prevalence of four different tail lesions in the slaughterhouse data correlated positively with annual mortality ( $r_s = 0.26, p = 0.02$ ) but not with the measures of condemnations. The percentage of tails with minor wounds in slaughterhouse correlated positively both with annual total carcass condemnations ( $r_s = 0.30, p = 0.007$ ) and with annual partial carcass condemnations ( $r_s = 0.26, p = 0.02$ ) but not with annual mortality. The percentage of tails with major wounds scored in the slaughterhouse did not correlate with any of the parameters of production parameters ( $r_s < 0.1$  for all).

**FIGURE 6 |** Percentage of undocked, tails with major acute wounds evaluated at slaughterhouse. The finishing pigs originated from 84 herds, where the herd veterinarian collected the data during one herd health visit. The results are shown separately for herds using wood as enrichment and according to explorative behavior of standing animals if they were not eating, drinking, defecating, or urinating (interaction), as evaluated by a veterinarian during the herd health visit. The behavior was directed toward enrichment material and not toward pen structures or pen mates. Different letters represent statistical difference:  $p < 0.001$ .

herds varied considerably and herds commonly had different from sudden-forceful or epidemic-type TB (30). Herds with a combinations of tail scorings evaluated at the slaughterhouse, high percentage of all types of lesions are the most problematic. Further, the correlations between the occurrence of healed tails, this indicates problems throughout the production period and acute tail lesions were rather weak. Therefore, against our need to improve intervention and preventive strategies when hypothesis, it is not possible to directly estimate the TL score outbreaks do occur. We suggest that by examining farm-wise the slaughterhouse by recording only one type of pigtail score lesion profiles it is possible to tailor advisory measures better (such as major acute wounds of the tail at the slaughterhouse) individual farms.

As recorded by a veterinarian during one herd health visit, the scoring at the slaughterhouse was performed by the tail score recorded at the slaughterhouse was only moderate researchers and was much more detailed than the one performed related to the on-farm TB level. We further observed some by veterinarians in the herds. However, as shown in Figures 3 and 4, farm evaluations on tail lesions and intact tails was somewhat related to the slaughterhouse evaluation of the tails. It is difficult to evaluate tail damage clinically in live pigs (31) since they are in groups in their pens. Dark tails of some breeds may be more difficult to evaluate on farm and especially tails in large farms might be more difficult to evaluate. At the herd level, it is not possible for evaluators to check each tail separately by manual palpation, which would be needed for a detailed classification, especially to identify healed damages. (One different herds, Figure 2) supports the notion that there are possibility is to examine tail posture in addition to lesions. I different types of TB (29). This also indicates that TB is likely a study investigating tail position, finishing pigs with wounds to occur at different stages of production in different farms. or in aimed wounds were 4.15 and 14.24 times more likely to Some herds had a high percentage of healed tails, with almost no acute wounds. This indicates that TB is a problem at an earlier stage of production, possibly in the weaning unit. Other (27, 32) did not include scoring of healed tails. Thus, we do not know if tail posture is a relevant measure of healed tail lesions. In our data, it is likely that veterinarians included both healed and healed tails in the same category while evaluating the tail became very severe. Further, some herds had a high percentage of addition in the herd, especially if the healed tails were not of major acute wounds, while only a moderate level of mild and drastically shortened and therefore the pigs were likely to take types of damage. This might indicate that the farms had suffered them up and curled. The average length of the healed tails in

slaughterhouse in this study was on average 97% of the length of the intact tails. Valros et al (26) suggested that healed tails with > 75% of the average intact length remaining could be considered intact enough.

Healed lesions are problematic from both an animal welfare and economical perspective. A healed tail has been bitten at some point and caused pain and stress in the pigs. No information is available about the time needed for the tail to heal after being bitten. The wound caused by tail docking heals in 4 to 8 weeks (33). After biting, the time needed for healing might be even longer as there is strong inflammation after TB (36). Healed tails might still have macroscopically invisible deep infections and their presence is associated with meat inspection findings and carcass condemnations at slaughterhouses. During the healing process, the tails are also likely to be promoting more TB than if the tails were totally intact. Furthermore, the timing of biting is difficult to evaluate after the tail has healed. Our fresh wood in addition to a straw rack, a metal chain, and slaughterhouse data show that the scores of both intact and healed tails are normally distributed. They are likely to describe the long-term situation both in the finishing herd and during earlier stages of the pig's life.

Meat inspection data could be better used in herd health and welfare planning at the farm level (34). Our results suggest that tail condition at the end of the finishing phase and long-term parameters of production quality (total and partial carcass condemnations and annual mortality) provide similar information about overall welfare of animals in the herd. It has previously been shown that tail lesions have been potential to be used as iceberg indicators of pig health and welfare (35). However, the scoring system probably affects the reliability of tail lesions as a measure of welfare. In the study by van Staaveren et al (34), tails were scored with a five-level scale, which is not performed as routine at the moment of slaughter. Keeling et al (26) also suggested that a three-level scale, including both old and new damage, should be used at slaughterhouse evaluations of the tails. However, based on our results, we carefully suggest that by evaluating the percentage of intact tails at the slaughterhouse it is possible to obtain a reasonable estimate of the long-term welfare situation of the farm.

Nearly half of the tails at the slaughterhouse were scored as intact, which is a very low figure. However, after dividing the tails into those without wounds (intact 48.1% and healed 37.3%, altogether 85%) and those with wounds (minor 12.4% and major acute wounds 2.3%, altogether 15%), the results seem to be in the same range with the results from other studies (21, 23, 29). However, comparison with the past studies is challenging as the data were collected from undocked or docked pigs and different scoring methods have been used. In addition, it is common to report the percentage of bitten tails and not the percentage of intact tails, which affects interpretation of results. Finally, the definition of an intact tail in this study is very strict, and as suggested by Valros et al (26), this level of precision is not feasible in a practical situation, such as at slaughterhouses would score tails on a continuous basis.

We showed that the results of different tail lesions at the slaughterhouse are associated with some herd-level parameters such as use of enrichment. EFSA (1) recommends that pigs should be given suitable materials to enable them to fulfil their needs to look for food, bite, root, and manipulate. The materials should be safe and at least one of the following qualities should be met: it should be edible/feed-like, chewable, investigate or manipulable. All our study farms provided their pigs with enrichment of some kind. Most farms provided hay, straw, or both, which are known to reduce the risk of TB on farms (36). However, the data did not allow for evaluation of the actual functionality of the enrichment. Although a recent review concluded that enrichment other than straw was not effective in TB prevention (37), the finding in the current study of the effect (tendency) of the using of wood was consistent with the study by Telkänranta et al (38). They found that provision of fresh wood in addition to a straw rack, a metal chain, and enrichment increased exploratory behavior while reducing ear biting when compared with pigs given no wood. In our study, wood as enrichment tended to be associated with both the percentage of intact tails and tails with major wounds. Notably, wood was especially associated with an increase in the percentage of intact tails in herds with leg problems other than arthritis. Further, wood appeared to have a protective role against major tail wounds on farms where explorative behavior was assessed to occur at a low level. This supports the finding by Telkänranta and Valros (39) that providing freshly harvested (but not dry) wood had an additive effect with straw in reducing tail-bite manipulation. Possibly, there is something in the texture of wood directing the innate need of the pigs to manipulate face other challenges, such as leg problems. However, we did not analyze the combinations of different enrichments or their amounts used, and therefore this finding should be interpreted with caution.

Sickness has been considered as a risk factor for TB (40). Two possible links between sickness and TB have been suggested (40). Recovered animals may have an increased propensity to become tail biters after illness or the increased attention of other pigs toward a sick one leads to an increased risk for the sick pig to become a victim of TB. The biggest health problems seemed to affect single individuals (usually 15%), the results of leg problems (arthritis, claw injuries, and other leg problems), presence of runt pigs or pigs with hernias, and skin injuries, infections, or abscesses, most of them likely to be associated with managerial problems. In univariate analyses, some of these conditions were associated with the tail scores, but in the final regression analysis only the presence of leg problems other than arthritis were associated with the percentage of healed tails. The association between disease conditions and tail lesions is difficult to assess in field studies, as data collected at herd level and at slaughterhouses. For example, Moirand

et al. (41) found an association between respiratory diseases, herds before slaughter. Herds seem to have varying combinations of rectal prolapse, and TB. The association between diseases and percentages of tails that were intact, healed, with minor TLs have been shown in studies using mostly pathological acute wounds, and with major acute wounds. Each of these slaughterhouse data. A connection has been found between provide different information about the situation on the farm. TL and abscesses (9, 26), arthritis or leg disorders (26, 31), pericarditis (26), respiratory conditions (7, 9, 23, 26, 31), skin infections (26), and whole and partial carcass health visit. Tail lesion scoring at the slaughterhouse appears to be a more precise measure than that performed on-farm as part of a herd condemnations (26). health visit. Tail condition as measured at the slaughterhouse shows potential to estimate the long-term welfare level on the farm. This is related to management, such as the use of manipulable objects, and maybe also to some health parameters on the farm.

It is difficult to find results about the possible association between mortality and TLs. Moinard et al. (41) reported that a post-weaning mortality of 2.5% was associated with a 3.9-fold increase in the risk of TB estimated on farms by researchers.

Mortality is a very rough measure to estimate animal welfare and the decisions of the farmer when to cull an animal has a large effect on this figure. However, the association between higher annual mortality and increase in healed tails in our study can be considered to describe the long-term situation of welfare of the herd. The association of increased percentage of intact tails and decreased mortality also supports the same idea of mortality being associated with tail score in the slaughterhouse.

Our study further shows that it is possible to grow undocked pigs with only a small percentage of pigs with serious tail wounds at the slaughterhouse. In almost 20% of the herds (Figure 1), no pigs had major acute wounds upon arrival at the slaughterhouse. Further, when examining the percentage of intact tails, at least 10% of the herds had 79% fully intact tails and the very best herd had 93% fully intact tails. Further, according to the herd health visit data, in 51.2% of the herds, veterinarians had recorded > 95% of tails as intact. This figure is well in accordance with the result of our scoring at the slaughterhouse, when we included both intact and healed tails in the figure.

This study has some limitations. It was not possible to evaluate the same pigs in the herd and at the slaughterhouse. However, we believe that the risk factors and situation in the herds are quite constant except for occasional TB outbreaks. The fact that piggeries usually empty their finishing units in 3–4 deliveries may have caused selection of certain types of pigs in our slaughterhouse data of 1 week. For example, the last delivery group of the pigs in one room may include those that have grown slower than the ones that were sent to slaughterhouse in the first delivery group. Due to these two factors, we may not have identified all real associations between the risk factors in the herd and the tail scoring results at slaughterhouse. However, we believe that the ones that have been identified are correct. The herd visits were performed by several different veterinarians, which also caused variation in the evaluation. However, this variation is likely mitigated by common educational background (veterinary studies and Sikava course).

## CONCLUSIONS

Our results show that by recording only one type of tail condition such as tails with major acute wounds at the slaughterhouse, it is not possible to estimate the total tail lesion situation

## DATA AVAILABILITY STATEMENT

The data of the slaughtered pigs are owned by the producers. Those interested in the datasets should contact Anna Valros (anna.valros@helsinki.fi). The Board of Sikava gave the permission to use the data anonymously.

## ETHICS STATEMENT

Ethical review and approval was not required for the animal study because the data was collected firstly from database where veterinarians had stored their herd health evaluation and secondly in the slaughterhouse after stunning and bleeding. Written informed consent for participation was not obtained from the owners because the data from the database was anonymous.

## AUTHOR CONTRIBUTIONS

AV was responsible for the data analysis. All authors contributed to the study design, data collection, interpretation of the results, preparation of the manuscript, participated in writing of the manuscript, and gave their approval for the final manuscript.

## FUNDING

The study was funded by HKScan Finland Oyj, the University of Helsinki, and the Kyllikki and Uolevi Lehtikainen fund. EF participation was possible through a short-term scientific mission (STSM reference number CA15134-44550) awarded by the COST ACTION GrouphouseNet (CA15134), which is supported by the EU Framework program Horizon 2020.

## ACKNOWLEDGMENTS

The authors acknowledge Riitta Pietilä and Johanna Daka for assistance in data collection at the slaughterhouse. The authors also acknowledge the abattoir personnel for their assistance during the data collection.



## REFERENCES

- Bracke MBM, De Lauwere CC, Wind SMM, Zonderland J. Attitudes of Dutch pig farmers towards tail biting and tail docking. *Agric Environ Ethics* (2013) 26:847–68. doi: 10.1007/s10806-012-9410-2
- D'Eath RB, Niemi JK, Vosough Ahmadi B, Rutherford KMD, Ison SH, Turner SP, et al. Why are most EU pigs tail docked? Economic and ethical analysis of four pig housing and management scenarios in light of EU legislation and animal welfare outcomes. *Animal*. (2016) 10:687–99. doi: 10.1017/S1751731115002098
- Heinonen M, Orro T, Kokkonen T, Munsterhjelm C, Peltoniemi O, Valros A. Tail biting induces a strong acute phase response and tail-end inflammation in nishings pigs. *Vet J Duxford* (2010) 184:303–7. doi: 10.1016/j.tvjl.2009.02.02
- Valros A, Munsterhjelm C, Puolanne E, Ruusunen M, Heinonen M, Peltoniemi OAT, et al. Physiological indicators of stress and meat carcass characteristics in tail bitten slaughter pigs. *Acta Vet Scand* (2013) 55:75. doi: 10.1186/1751-0147-55-75
- Munsterhjelm C, Brunberg E, Heinonen M, Keeling L, Valros A. Stress measures in tail biters and bitten pigs in a matched case-control study. *Anim Welfare* (2013) 22:331–8. doi: 10.7120/09627286.22.3.331
- Valros A, Ahlström S, Rintala H, Häkkinen T, Saloniemi H. The prevalence of tail damage in slaughter pigs in Finland and association to carcass condemnation. *Acta Agric Scand A An Sc* (2004) 54:213–9. doi: 10.1080/09064700510009234
- Kritas SK, Morrison RB. Relationships between tail biting and disease lesions and condemnations at slaughter. *Met. Rec.* (2007) 160:149–52. doi: 10.1136/vr.160.5.149
- Harley S, More SJ, Boyle L, O'Connell NE, Hanlon A. Good animal welfare makes economic sense: potential pig abattoir meat inspection as a welfare surveillance tool. *Irish Vet J* (2012) 65:11. doi: 10.1186/2046-0481-65-11
- vom Brocke AL, Karnholz C, Madey-Rindermann D, Gauly M, Leeb C, Winckler C, et al. Tail lesions in fattening pigs: relationships with postmortem meat inspection and in uence of a tail biting management protocol. *Animal*. (2018) 13:835–44. doi: 10.1017/S1751731118002070
- Sinisalo A, Niemi J, Heinonen M, Valros A. Tail biting and production performance in fattening pigs. *Livest Sci.* (2012) 143:142–50. doi: 10.1016/j.livsci.2011.09.019
- EFSA Panel on Animal Health and Welfare. EFSA (European Food Safety Authority) Scientific Opinion concerning a multifactorial approach in the use of animal and non-animal-based measures to assess the welfare of pigs. *EFSA J* (2014) 12:3702. doi: 10.2903/j.efsa.2014.3702
- Taylor NR, Parker RMA, Mendl M, Edwards SA, Main DCJ. Prevalence and risk factors for tail biting on commercial farms and intervention strategies. *Vet J*. (2012) 194:77–83. doi: 10.1016/j.tvjl.2012.03.004
- Lahrmann HP, Busch ME, D'Eath RB, Forkman B, Hansen CF. More tail lesions among undocked than tail docked pigs in a conventional system. *Animal*. (2017) 11:1825–31. doi: 10.1017/S1751731117000490
- Kongsted H, Foldager L, Sorensen JT. Data from routine meat inspection is a poor indicator of the prevalence of tail lesions in undocked pigs. *Porcine Health Manag* (2020) 6:10. doi: 10.1186/s40813-020-00149-z
- Calderon Diaz JA, Boyle LA, Diana A, Leonard FC, Moriarty JP, Harte MC, et al. Early life indicators predict mortality, illness, reduced welfare and carcass characteristics in nisher pigs. *Prev Vet Med* (2017) 146:94–102. doi: 10.1016/j.prevetmed.2017.07.018
- Carroll GA, Boyle LA, Hanlon A, Collins L, Gri n K, Friel M, et al. What can carcass-based assessments tell us about the lifetime welfare of pigs? *Livestock Sci* (2018) 214:98–105. doi: 10.1016/j.livsci.2018.04.020
- Pandolf F, Edwards SA, Maes D, Kyriazakis I. Connecting real data sources to assess the interconnections between biosecurity, health, welfare, and performance in commercial pig farms in Great Britain. *Front Vet Sci*. (2018) 5:41. doi: 10.3389/fvets.2018.00041
- van Staaveren N, Doyle B, Manzanilla EG, Calderón Díaz JA, Hanlon A, Boyle L. Validation of carcass lesions as indicators for on-farm health and welfare of pigs. *J Anim Sci* (2017) 95:1528–36. doi: 10.2527/jas.2016.1180
- Valros A, Heinonen M. Save the pig tail. *Porcine Health Manag* (2015) 1:2. doi: 10.1186/2055-5660-1-2
- Carroll CA, Boyle LA, Teixeira DL, van Staaveren N, Hanlon A, O'Connell NE. Effects of scalding and dehairing of pig carcasses at abattoirs on the visibility of welfare-related lesions. *Animal*. (2016) 10:460–7. doi: 10.1017/S1751731115002037
- Harley S, More SJ, O'Connell NE, Hanlon A, Teixeira DL, Boyle LA. Evaluating the prevalence of tail biting and carcass condemnation in slaughter pigs in the Republic and Northern Ireland, and the potential abattoir meat inspection as a welfare surveillance tool. *Met. Rec.* (2012) 171:621. doi: 10.1136/vr.100986
- Harley S, Boyle L, O'Connell NE, More SJ, Teixeira DL, Hanlon A. Docking the value of pigmeat? Prevalence and nancial implications of welfare lesions in Irish slaughter pigs. *Anim Welfare*. (2014) 23:275–85. doi: 10.7120/09627286.23.3.275
- van Staaveren N, Vale AP, Manzanilla EG, Teixeira DL, Leonard FC, Hanlon A, et al. Relationship between tail lesions and lung health in slaughter pigs. *Prev Vet Med* (2016) 127:21–6. doi: 10.1016/j.prevetmed.2016.03.004
- Lemos Teixeira DL, Harley S, Hanlon A, O'Connell NE, More SJ, Manzanilla EG, et al. Study on the association between tail lesion score, carcass weight, and viscera condemnations in Slaughter pigs. *Front Vet Sci* (2016) 3:24. doi: 10.3389/fvets.2016.00024
- Keeling LJ, Wallenbeck A, Larsen A. Scoring tail damage in pigs: evaluation based on recordings at Swedish slaughterhouses. *Acta Vet Scand* (2012) 54:32. doi: 10.1186/1751-0147-54-32
- Valros A, Välimäki E, Nordgren H, Vugts J, Fàbrega E, Heinonen M, et al. Tail lesions as a welfare indicator in nishing pigs? Scoring of tail lesions and de nting intact tails in undocked pigs at the abattoir. *Front Vet Sci* (2020) 7:405. doi: 10.3389/fvets.2020.00405
- Wallgren T, Larsen A, Gunnarsson S. Tail posture as an indicator of tail biting in undocked nishing pigs. *Animals*. (2019) 9:18. doi: 10.3390/ani9010018
- Pig subgroup of the EU platform on animal welfare. Output of the Commission's Subgroups on Transport & P (2019). Available online at: [https://ec.europa.eu/food/sites/food/files/animals/docs/platform\\_20191007\\_pres-03.p](https://ec.europa.eu/food/sites/food/files/animals/docs/platform_20191007_pres-03.p) (accessed December 7, 2020).
- Taylor NR, Main DCJ, Mendl M, Edwards SA. Tail-biting: a new perspective. *Vet J* (2010) 186:137–47. doi: 10.1016/j.tvjl.2009.08.028
- Valros A. Tail biting. In: Spinka M, editor. *Advances in Pig Welfare*, 2nd edn. Woodhead Publishing, Elsevier, UK (2018). p. 137–66.
- Munsterhjelm C, Simola O, Keeling L, Valros A, Heinonen M. Health parameters in tail biters and bitten pigs in a case-control study. *Animal*. (2013) 5:814–21. doi: 10.1017/S1751731112002194
- Zonderland JJ, van Riel JW, Bracke MBM, Kemp B, Den Hartog LA, van Soest HAM. Tail posture predicts tail damage among weaned piglets. *Appl Anim Behav Sci* (2009) 121:165–70. doi: 10.1016/j.applanim.2009.09.002
- Sandercock DA, Smith SH, Di Gimignano P, Edwards SA. Histopathological characterization of tail injury and traumatic ulcer development after tail docking in piglets. *J Comp Pathol*. (2016) 155:40–9. doi: 10.1016/j.jcpa.2016.05.003
- Devitt C, Boyle L, Teixeira DL, O'Connell NE, Hawe M, Hanlon A. Pig producer perspectives on the use of meat inspection as an animal health and welfare diagnostic tool in the Republic of Ireland and Northern Ireland. *Irish Vet J* (2016) 69:2. doi: 10.1186/s13620-016-0076-3
- Van Staaveren N, Teixeira DL, Hanlon A, Boyle L. Pig carcass lesions: the in uence of record keeping through an advisory service and the relationship with farm performance parameters. *Animal*. (2017) 11:140–6. doi: 10.1017/S1751731116001117
- Wallgren T, Larsen A, Lundeheim N, Westin R, Gunnarsson S. Condition and impact of straw provision on behaviour, lesions and pen hygiene of commercial farms rearing undocked piglets. *Appl Anim Behav Sci* (2019) 210:26–37. doi: 10.1016/j.applanim.2018.10.013
- Buijs, S, Muns R. A review of the effects of non-straw enrichment on tail biting in pigs. *Animals*. (2019) 9:824. doi: 10.3390/ani9100824
- Telkänranta H, Bracke MBM, Valros A. Fresh wood reduces tail biting and increases exploratory behaviour in nishing pigs. *Appl Anim Behav Sci*. (2014) 161:51–9. doi: 10.1016/j.applanim.2014.09.007
- Telkänranta H, Valros A. Pigs with but not without access to fresh wood of recently harvested wood show reduced pen-mate manipulation after a provision of feed and straw. *Appl Anim Behav Sci* (2020) 232:105103. doi: 10.1016/j.applanim.2020.105103
- Munsterhjelm C, Nordgreen J, Aae F, Heinonen M, Valros A, Jancza A. Sick and grumpy: changes in social behaviour after a controlled

- immune stimulation in group-housed gilts *Physiol Behav* (2019) 198:76–83. doi: 10.1016/j.physbeh.2018.09.018
41. Moinard C, Mendl M, Nicol CJ, Green LE. A case control study of farm risk factors for tail biting in pigs *Appl Anim Behav Sci* (2003) 81:333–55. doi: 10.1016/S0168-1591(02)00276-9
- Conflict of Interest: EV and JV are employees of the slaughterhouse company. The University of Helsinki was responsible for data analysis and scientific publication throughout the study. The company involvement did not have any effect on the study design, data collection, interpretation of results, or writing of the manuscript.
- The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Copyright © 2021 Heinonen, Välimäki, Laakkonen, Toppari, Varga and Valros. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.